

Common Name of Parasites

Dr. Anita Rathore

Assistant Professor, Department of Veterinary Pathology, College of Veterinary and Animal science, Navania, Vallabh Nagar, Udaipur
<https://doi.org/10.5281/zenodo.8359402>

1. *Acanthocephalan* – thorny headed worm
2. *Ancylostoma duodenale* – hook worm (man)
3. *Ancylostoma tubaeforme* – hook worm (cat)
4. *Anoplocephala magna* – large horse worm
5. *Anoplocephala perfoliata* – lappet tapeworm
6. *Binitellobitharzia* – portal vein of elephant
7. *Bunostomum phlebotomum* - hook worm.
8. *Bunostomum trigonocephalum* - hook worm (sheep & goat)
9. *Capillaria species* – bobilo
10. *Chlonorchis sinensis*
11. *Chlonorchis sinensis* or *Oopistherchis sinensis* – oriental or Chinese liver fluke
12. *Corynosoma strumosum* – thorny headed worm
13. *Cotugnia digonopora* – bipore tapeworm of poultry
14. *Cutaneous habronemiasis* – other name is summer sore , bersati, granular dermatitis
15. *Davainea proglottina* – smallest poultry tapeworm but most pathogenic
16. *Davaineidae family* – hammer shaped hooks
17. *Dicrocoelium dentriticum* – lancet fluke
18. *Dictyocaulus filaria* – boot shape spicule, milky white
19. *Dilepididae family* – rose thorn shaped hooks
20. *Diectophyma renale* – largest nematode – kidney worm of dog
21. *Diphyllobothrium latum* – public health importance – macrocytic hypochromic anemia
22. *Dipylidium caninum* – commonest and bipolar tapeworm of dog
23. *Dirofilaria immitis* – dog heart worm
24. *Draschia megastoma* – nodule form in stomach
25. *Draschia megastoma* and *Habronema muscae* – musca domestica
26. *Echinococcus granulosus* – zoonotic tapeworm
27. *Echinostomatidae* – head collar present
28. *Enterobius vermicularis* – pin worm or seat worm in man
29. *Fasciolopsis buskii* – raw water chestnut (eating) (transmitted)
30. *Filicolis anatis* – thorny headed worm

31. *Fimbriaria fasciolaris* – pseudo scolex
32. *Gaigeria pachyscelius* – hookworm (sheep & goat)
33. *Gastrodiscus agyptiacus* – saucers shaped body
34. *Gigantobitharzia* – birds
35. *Habronema majus* – stable fly (*Stomoxys calcitrans*)
36. *Haemonchus contortus* – wire worm, barbers pole worm, twisted worm, stomach worm, dorsal lancet, found in abomasum of sheep
37. *Heterakis gallinarum* - caecal worm of poultry
38. *Heterophidae* family – smallest trematodes
39. *Hydatid cyst* – metacestodes – larval stage
40. *Hymenolepis nana* – dwarf tapeworm of human being and rodents, autoinfection
41. *Macrocanthorhynchus hirudinacius* – thorny headed
42. *Micronema detetrii* – cerebrospinal nematodes (horse)
43. *Moniezia benedeni* – double pore
44. *Moniezia expansa* – largest tapeworm (double pore)
45. *Muellerius capillaris* – lung worm of sheep (commonest)
46. *Necator americanus* - hook worm (man)
47. *Nematomorphs* – hair worm
48. *Neorickettsia helminthrica* – solomon poisoning
49. *O. equi*- false whip worm
50. *Oesophagostomum radiatum* - pimply gut-found in intestine of cattle
51. *Oncicola canis* – thorny headed
52. *Opisthorchis fecinens*
53. *Opisthorchis sinensis*
54. *Opisthorchis tenuicollis*
55. *Oxyuris equi* – false whip worm or horse pin worm, single pin shape spicule
56. *Paramphistomatidae* family – plug feeder
57. *Paramphistomum cervi* – conical flukes
58. *Paranoplocephala mamillana* – without lappet and dwarf tapeworm
59. *Parascaris equorum* – large round worm of horse
60. *Polymorphus boschadis* – thorny headed worm
61. *Prosthogonimidae* – grape like bunches
62. *Raillietina echinobothrida* – nodular tapeworm
63. *Raillietina tetragona* – largest fowl worm
64. *Setaria cervi* – buffalo
65. *Setaria digitata* – cattle and buffalo
66. *Setaria labiatopapillosa* – sheep, goat
67. *Setaria species*- eye worm (cerebrospinal nematodiasis)
68. *Stephanofilaria assamensis* – Hump sore
69. *Stephanofilaria zaheeri* - ear sore
70. *Stephanurus dentatus* - kidney worm of swine
71. *Stilesia hepatica* – dumbbell shaped uterus
72. *Strongylus edentates* - palisade worm
73. *Strongylus equi* - palisade worm
74. *Strongylus vulgaris* – palisade worm
75. *Subulura brumpti* - caecal worm of poultry



76. *Syngamus trachea* - gapeworm of poultry
77. *Taenia multiceps* – gid or staggers (brain)
78. *Taenia taeniaformis* – bell shaped, post proglottis, strobilocercus fasciolaris
79. *Thelazia alfortensis* – cattle
80. *Thelazia californionsis* – sheep eye worm
81. *Thelazia gulosa* – cattle eye worm
82. *Thelazia lacrymalis* - eye worm of horse
83. *Thelazia rhodesia* - (common in all) - ruminant eye worm
84. *Thysaniezia giardia* – undulating uterus
85. *Thysaniezia helictometra*
86. *Thysanosoma actinoides* – fringed tapeworm
87. *Toxocara canis* - arrow worm of dog and canine
88. *Toxocara cati* – arrow worm of cat and feline
89. *Toxocara leolina* - arrow headed worm (cervical alae present)
90. *Toxocara transfuga* - arrow worm
91. *Toxoplasma* and *Sarcocystis* - endodyogeny (protozoans)
92. *Trichinella spiralis* – Smallest nematode
93. *Trichostrongylus axei* - Ring worm
94. *Trichuris globulosa* – Ceccal worm of camel
95. *Trichuris ovis* – whip worm, longer male than female
96. *Uncinata stenocephala* - hook worm of dog

