THE SCIENCE WORLD
A Monthly e Magazine
ISSN:2583-2212
Popular Article
August, 2026 Vol.6(8), 2507-2512

Aparajita: The Blue Blossom of Nature

Tanmeet Singh[1]*, Julie Dogra Bandral[1], Monika Sood[1], Jesreen Chauhan[1] and Arushi Sharma[1]
Division of Post Harvest Management
Sher-e-Kashmir University of Agricultural Sciences and Technology of Jammu
***Corresponding Author-Email:** singhtanmeet18@gmail.com
https://doi.org/10.5281/zenodo.23469678

Abstract

Clitoria ternatea Linn., commonly known as **Aparajita** or **butterfly pea**, is a perennial herbaceous climbing plant belonging to the family **Fabaceae**. It is widely cultivated as an ornamental vine because of its attractive blue or white flowers and has recently gained considerable attention in the food industry as a natural source of blue anthocyanin pigments. The plant is a slender, twining climber that can grow up to 2–3 m in height when supported. It possesses green, branched stems, alternate leaves, and a well-developed taproot with nitrogen-fixing nodules, enabling it to improve soil fertility. Its large, butterfly-shaped flowers, characteristic of the pea family, are predominantly deep blue with a white or yellow throat, although pink and violet variants are also found. Beyond its ornamental value, *C. ternatea* has been recognized for its diverse medicinal properties, including antifungal, antidiabetic, and anti-inflammatory activities.

The plant produces flat, linear pods measuring approximately 5–7 cm in length, each containing seven to ten smooth seeds that range from yellowish-brown to black in colour. Young pods are edible and consumed as vegetables in some regions. Propagation occurs mainly through seeds, which germinate within one to two weeks under warm conditions, while flowering generally begins about four weeks after germination. Owing to its rapid growth, drought tolerance, and ability to thrive in marginal soils, *C. ternatea* has become an economically important species for both sustainable agriculture and environmental conservation (Jeyaraj *et al*., 2021).

Butterfly pea flowers are particularly valued for their intense blue pigmentation, which is produced through anthocyanin biosynthesis involving enzymes such as **chalcone synthase (CHS), chalcone isomerase (CHI), flavanone-3-hydroxylase (F3H), and anthocyanin synthase (ANS)**. Even small quantities of the flowers yield a vibrant natural colour, making

them suitable for food applications. Among their most popular uses is the preparation of **blue tea**, which is often blended with spices such as ginger, cardamom, and cinnamon to enhance both flavour and health-promoting properties. In traditional Ayurvedic medicine, Aparajita has long been regarded as a **Medhya Rasayana** (brain tonic) and has been used to improve memory, relieve stress, treat skin disorders, and manage several chronic ailments. Contemporary scientific research has further confirmed its rich phytochemical profile and numerous biological activities, including antioxidant, antimicrobial, anti-inflammatory, antidiabetic, neuroprotective, and anticancer effects (Tripathi *et al*., 2022).

Originally believed to have originated in tropical Asia, *C. ternatea* is now widely distributed across India, Southeast Asia, Australia, Africa, and parts of Central and South America. Its adaptability allows it to flourish under diverse environmental conditions, particularly in warm, humid climates with fertile, well-drained soils and adequate sunlight. Because of its vigorous growth and low maintenance requirements, it is extensively cultivated in home gardens, agricultural fields, and landscaped areas (Jeyaraj *et al*., 2021).

Apart from its ornamental significance, butterfly pea flowers have gained substantial commercial value as natural colourants for beverages, confectionery, bakery products, dairy items, desserts, and herbal teas. The anthocyanins present in the petals exhibit pH-dependent colour changes, making them suitable for the development of functional foods and intelligent food packaging systems. Unlike synthetic food dyes, these pigments are biodegradable, non-toxic, and possess strong antioxidant properties, thereby meeting the increasing consumer demand for safe and natural food ingredients (Jeyaraj *et al*., 2021).

Phytochemical investigations have revealed that *C. ternatea* is rich in tannins, flavonoids, glycosides, steroids, triterpenoids, phenolic compounds, and saponins, all of which contribute to its diverse pharmacological activities. The extraction efficiency of these bioactive compounds is influenced by factors such as solvent type, extraction temperature, pH, and extraction technique. Consequently, advanced methods including **ultrasound-assisted extraction** and **optimized aqueous ethanol extraction** have been developed to improve the recovery of anthocyanins and phenolic compounds for applications in the food, nutraceutical, and pharmaceutical industries (Tripathi *et al*., 2022).

Phytochemical Composition and Therapeutic Significance

The therapeutic value of *Clitoria ternatea* is largely associated with its rich and diverse phytochemical profile. The plant contains a wide range of biologically active compounds, including anthocyanins, flavonoids, flavanol glycosides, phenolic acids, tannins, saponins, triterpenoids, steroids, phytosterols, and tocopherols. Among these, the flowers are particularly

Official Website
www.thescienceworld.net
thescienceworldmagazine@gmail.com
Published 30/08/2026

rich in **ternatins**, a distinctive group of delphinidin-derived anthocyanins responsible for the characteristic deep blue colour of butterfly pea flowers. Besides imparting colour, ternatins exhibit excellent pigment stability and potent antioxidant activity. Other important phytochemicals, such as kaempferol, quercetin, and myricetin glycosides, along with various phenolic compounds, further enhance the medicinal and functional properties of the plant. Phytochemical analyses have also confirmed the occurrence of glycosides, tannins, terpenoids, and saponins in flower extracts, highlighting *C. ternatea* as a valuable natural reservoir of bioactive secondary metabolites (Tripathi *et al*., 2022).

The quantity and quality of these bioactive compounds are greatly influenced by the extraction method employed. Conventional techniques, including aqueous extraction, ethanol extraction, and maceration, have been extensively used to isolate phytochemicals from butterfly pea flowers. However, recent research has demonstrated that advanced extraction methods, particularly **ultrasound-assisted extraction** and **optimized aqueous ethanol extraction**, significantly improve the recovery of anthocyanins, flavonoids, and phenolic compounds. Furthermore, maintaining a lower extraction pH and moderate processing temperatures enhances anthocyanin stability by preserving the pigments in their flavylium cation form, thereby reducing degradation and increasing extraction efficiency. These optimized extraction strategies have expanded the utilization of butterfly pea flower extracts in the formulation of functional foods, nutraceuticals, and natural health products (Jeyaraj *et al*., 2021).

Health Benefits and Agricultural Importance of *Clitoria ternatea*

1. Potent Antioxidant Activity

One of the most widely investigated biological properties of *Clitoria ternatea* is its strong antioxidant potential. Excessive generation of reactive oxygen species (ROS) leads to oxidative stress, which plays a significant role in the onset and progression of chronic diseases such as cardiovascular disorders, diabetes, neurodegenerative diseases, and cancer. The flower is rich in anthocyanins and phenolic compounds that effectively scavenge free radicals by donating electrons or hydrogen atoms, thereby protecting lipids, proteins, and DNA from oxidative damage. Experimental studies have demonstrated that butterfly pea flower extracts possess excellent free radical scavenging capacity and inhibit lipid peroxidation, making them valuable natural antioxidants for use in functional foods, nutraceuticals, and dietary supplements (Tripathi *et al*., 2022).

Official Website
www.thescienceworld.net
thescienceworldmagazine@gmail.com
Published 30/08/2026

2. Antidiabetic Potential

Clitoria ternatea has also attracted considerable attention for its antihyperglycaemic properties. Diabetes mellitus is characterized by prolonged elevation of blood glucose levels resulting from impaired insulin secretion or insulin resistance, often leading to complications affecting the cardiovascular, renal, nervous, and visual systems. Research indicates that flavonoids and anthocyanins present in butterfly pea flowers inhibit carbohydrate-digesting enzymes such as **α-amylase** and **α-glucosidase**, thereby slowing glucose absorption and reducing postprandial blood sugar levels. Experimental studies using diabetic animal models have reported significant reductions in fasting blood glucose, glycated haemoglobin (HbA1c), and malondialdehyde (MDA) levels following administration of flower extracts. These findings suggest that the plant may help improve glycaemic control while reducing oxidative stress and delaying diabetes-related complications through its antioxidant and antiglycation activities (Dewi *et al*., 2023).

3. Antimicrobial and Anti-inflammatory Properties

Butterfly pea has demonstrated broad-spectrum antimicrobial activity against several pathogenic microorganisms, making it a promising natural alternative to synthetic antimicrobial agents. Extracts obtained from different plant parts, including flowers, leaves, stems, and roots, have shown inhibitory effects against bacteria such as *Staphylococcus aureus, Escherichia coli, Bacillus cereus, Pseudomonas aeruginosa, Klebsiella pneumoniae, Proteus vulgaris, Salmonella typhi,* and *Aeromonas hydrophila*. This antimicrobial activity is largely attributed to the presence of flavonoids, phenolic compounds, alkaloids, terpenoids, and other secondary metabolites that disrupt microbial cell membranes and metabolic functions. Consequently, *C. ternatea* has potential applications in both pharmaceutical formulations and natural food preservation systems (Jamil *et al*., 2018).

In addition to its antimicrobial effects, the plant exhibits significant anti-inflammatory activity. Chronic inflammation is associated with numerous disorders, including arthritis, cardiovascular diseases, and metabolic syndromes. Bioactive compounds such as flavonoids and phenolics help suppress inflammatory mediators, regulate pro-inflammatory signalling pathways, and reduce oxidative stress-induced tissue damage. These scientific findings support the long-standing Ayurvedic use of Aparajita for managing inflammatory conditions and promoting overall health (Jeyaraj *et al*., 2021).

4. Emerging Anticancer Potential

Recent investigations have highlighted the promising anticancer properties of butterfly pea flower extracts. Since oxidative stress, persistent inflammation, and uncontrolled cell

Official Website
www.thescienceworld.net
thescienceworldmagazine@gmail.com
Published 30/08/2026

proliferation contribute to cancer development, the antioxidant-rich phytochemicals present in *C. ternatea* have gained considerable research interest. Anthocyanins, flavonoids, and phenolic compounds have demonstrated antiproliferative effects by suppressing tumour cell growth and inducing programmed cell death in experimental models. Although current evidence is primarily based on laboratory studies, these findings suggest that *C. ternatea* may serve as a valuable source of bioactive compounds for future anticancer drug development. Nevertheless, further preclinical and clinical investigations are required before its therapeutic applications can be fully established (Jeyaraj *et al*., 2021).

5. Role in Sustainable Agriculture

Apart from its medicinal significance, *Clitoria ternatea* also contributes substantially to sustainable agricultural systems. As a leguminous species, it develops a deep taproot system with numerous lateral roots that establish symbiotic associations with *Rhizobium* bacteria. These microorganisms fix atmospheric nitrogen into plant-available forms, thereby enhancing soil fertility and reducing the need for synthetic nitrogen fertilizers. Owing to this characteristic, butterfly pea is widely utilized as a **cover crop, green manure, and forage crop**, where it improves soil health, suppresses weed growth, and supports environmentally sustainable farming practices (Jeyaraj *et al*., 2021).

Clitoria ternatea L. (Aparajita) has emerged as a highly valuable medicinal and functional plant owing to its rich phytochemical composition, diverse therapeutic properties, and extensive applications across the food, pharmaceutical, and agricultural sectors. Revered in traditional Ayurvedic medicine as a **Medhya Rasayana**, it has been used for centuries to enhance cognitive function, alleviate inflammatory conditions, promote skin health, and support the management of various chronic ailments. Modern scientific investigations have provided substantial evidence supporting these traditional uses by demonstrating the plant's potent antioxidant, antidiabetic, antimicrobial, anti-inflammatory, neuroprotective, and anticancer activities, which are largely attributed to its abundance of anthocyanins, flavonoids, and phenolic compounds (Jeyaraj *et al*., 2021).

The flowers of *C. ternatea* are among the richest natural sources of **ternatins**, distinctive blue anthocyanin pigments that not only impart an attractive natural colour but also possess remarkable antioxidant properties. Their excellent colour stability and pH-sensitive characteristics have led to their increasing use as natural food colourants, functional food ingredients, and intelligent packaging indicators, thereby expanding their commercial significance in the food, nutraceutical, and pharmaceutical industries. In addition to its medicinal value, the plant contributes to sustainable agriculture through biological nitrogen

Official Website
www.thescienceworld.net
thescienceworldmagazine@gmail.com
Published 30/08/2026

fixation, improving soil fertility and serving effectively as a forage crop, green manure, and soil conditioner (Tripathi *et al*., 2022).

Despite the considerable progress achieved in understanding the biological and pharmacological potential of *Clitoria ternatea*, further clinical research is required to establish standardized dosages, confirm long-term safety, and validate its therapeutic efficacy in humans. Future studies should also focus on developing more efficient extraction techniques, enhancing anthocyanin stability, and exploring innovative applications in functional foods, pharmaceutical formulations, and intelligent food packaging systems. Overall, Aparajita represents a promising natural resource whose traditional medicinal value, supported by contemporary scientific evidence, underscores its significant potential to promote human health, drive innovation in natural product development, and contribute to sustainable agricultural practices.

References

Dewi, I., Chodidjah, C. and Atina, H. (2023) 'Evaluation of Clitoria ternatea L. flower extract in preventing complications of diabetes mellitus', Tropical Journal of Natural Product Research, **7**(10): 4908–4911.

Jamil, N. and Pa'ee, F. (2018) 'Antimicrobial activity from leaf, flower, stem and root of Clitoria ternatea: A review', AIP Conference Proceedings, 2002, 020044.

Jeyaraj, E.J., Lim, Y.Y. and Choo, W.S. (2021) 'Extraction methods of butterfly pea (Clitoria ternatea) flower and biological activities of its phytochemicals', Journal of Food Science and Technology, **58**(6) : 2054–2067.

Tripathi, R., Baghel, S.B.S., Hingwasiya, S. and Upadhyay, R. (2022) 'Qualitative analysis of Clitoria ternatea Linn. (Aparajita) flower of family Fabaceae', Indian Journal of Applied and Pure Biology, **37**(1) :110–114.