THE SCIENCE WORLD
A Monthly e Magazine
ISSN:2583-2212
Popular Article
March, 2026 Vol.6(3), 810-814

Bovine Lameness: A Modern Perspective on Prevention and Management

[1]Chithra S., [1]Nandhana Vijayakumar, [1]Shan Muhammed., [1] Alanis Lilly Cubeleo and [2]N. S. Sunilkumar
[1]BVSc&AH graduate students
[2]Assistant Professor, Department of Veterinary Anatomy
College of Veterinary and Animal Sciences, Mannuthy, Thrissur, Kerala, India
Kerala Veterinary and Animal Sciences University
doi.org/10.5281/ScienceWorld.19496853

Introduction

Bovine lameness is defined as an abnormality in a cow's movement or posture that is caused by pain, usually in the feet, legs or joints. Far from being a simple ailment, it is a complex, multifactorial disease that a significant challenge to cattle welfare and farm profitability worldwide. While lameness has long been recognized as a serious issue, our understanding of its underlying causes and pathology has undergone a significant paradigm shift. The focus has moved from a broad, general concept of hoof inflammation to a more precise understanding of specific, mechanically-induced lesions. This revised perspective is critical for developing effective prevention and treatment strategies that address the root causes of the problem.

Etiology: From Sub Acute Ruminal Acidosis (SARA) to Claw Horn Disruption Lesions (CHDL)

Historically, lameness was often linked directly to metabolic events like subacute ruminal acidosis (SARA). The theory was that a low rumen pH caused by high-starch diets led to the release of endotoxins, triggering systemic inflammation and compromising the hoof's structure. While SARA and nutritional imbalances are still recognized as contributing factors, the current understanding of lameness in cattle is centered on a more specific pathology: **Claw Horn Disruption Lesions (CHDL)**.

CHDL is a collective term for non-infectious hoof lesions, including sole ulcers, white line disease and sole hemorrhages. These lesions are now understood to be the result of a

cascade of events that culminates in mechanical disruption of the hoof horn-producing tissues (the corium). The true pathogenesis is a combination of nutritional, physiological and environmental stressors that weaken the hoof's internal structures and predispose it to trauma.

The Critical Role of the Digital Cushion

The **digital cushion** is a vital anatomical structure located beneath the pedal bone within the hoof capsule. Composed of a specialized mix of fat and fibrous tissue, its primary function is to act as a natural shock absorber, protecting the delicate corium and underlying bone from the concussive forces of walking. The structural integrity of the digital cushion is now considered a key determinant of hoof health.

Research has shown that the digital cushion's thickness and volume are directly related to a cow's body condition. During the periparturient period, when cows are in a state of negative energy balance and mobilise fat stores, the digital cushion can thin out. A thinner digital cushion offers less protection, leaving the corium vulnerable to compression and trauma from hard surfaces. This trauma leads to hemorrhage and a localized interruption of horn production, ultimately resulting in the formation of CHDLs. Therefore, maintaining a healthy body condition, especially around calving, is crucial for preserving the integrity of this internal protective mechanism and preventing lameness.

Clinical Signs and Symptoms

The clinical presentation of bovine lameness can vary in severity but often follows a predictable pattern of changes in gait, posture and behavior. The earliest and most subtle signs can include a slight shift in weight bearing or a reluctance to put full weight on one limb. As the condition progresses, more obvious symptoms emerge:

- **Changes in Locomotion:** The cow may walk with a noticeable limp, take short, hesitant steps, or show a reduced stride length. Head bobbing is a classic sign where the cow's head rises when the painful limb touches the ground to compensate for the discomfort.
- **Altered Posture:** A hallmark sign of lameness is an arched back, particularly when the cow is walking. In severe cases, this arched posture may persist even when the animal is standing still as it attempts to offload weight from the affected foot.
- **Behavioral Indicators:** Lame cows often show a reluctance to move, preferring to lie down for extended periods. They may be slow to rise, eat less and spend less time at the feed bunk, all of which contribute to reduced milk production and body condition loss.

Official Website
www.thescienceworld.net
thescienceworldmagazine@gmail.com
Published 30/03/2026

Diagnosis

Timely and accurate diagnosis is essential for effective treatment and long-term control of lameness. It requires a multi-step approach that combines careful observation with a thorough physical examination.

1. **Locomotion Scoring:** The first step is to systematically assess the entire herd. Locomotion scoring systems, such as the widely used 5-point scale, provide a standardized method for classifying the severity of lameness based on a cow's gait. Regular scoring allows for the early detection of subclinical cases before they become severe.
2. **Clinical Examination:** Once a lame cow is identified, a detailed clinical exam of the foot is necessary. This involves cleaning the hoof to remove debris, visually inspecting for swelling or lesions, and using specialized hoof testers to pinpoint areas of pain. Lifting and examining the hoof's sole and white line is crucial for identifying underlying issues like sole ulcers or abscesses.
3. **Advanced Diagnostic Tools:** In complex cases, advanced tools can provide further insight. Radiography can be used to identify fractures or deep bone infections, while ultrasonography may reveal soft tissue damage. These tools, while not routine, are invaluable for confirming difficult diagnoses.

Treatment

Effective treatment of bovine lameness hinges on a prompt, accurate diagnosis and a targeted therapeutic approach. The primary goals are to alleviate pain, heal the lesion and restore normal function.

1. **Pain Management:** Pain is a significant welfare issue and an impediment to recovery. The use of non-steroidal anti-inflammatory drugs (NSAIDs) such as flunixin meglumine or meloxicam is a cornerstone of treatment. NSAIDs reduce inflammation and pain, encouraging the cow to move more naturally and facilitating the healing process.
2. **Therapeutic Hoof Trimming:** Corrective hoof trimming is a critical step for most hoof-related lameness. A skilled trimmer can redistribute weight away from the painful area, remove compromised horn tissue and expose abscesses for drainage. This procedure is designed to provide immediate relief and promote healing of the corium.
3. **Hoof Blocks:** When a lesion is severe, a hoof block is often applied to the healthy claw. This elevates the affected claw off the ground, effectively transferring the

Official Website
www.thescienceworld.net
thescienceworldmagazine@gmail.com
Published 30/03/2026

animal's weight to the sound one. The affected claw is then free from pressure, allowing it to heal without further trauma.

4. **Topical and Systemic Treatments:** For infectious conditions like digital dermatitis or foot rot, topical antiseptics and antibiotics are used. In more severe systemic infections, a course of systemic antibiotics is administered under veterinary guidance.

Prevention: An Integrated Approach

While treatment is necessary, a proactive approach focused on prevention is far more effective and economical. A successful lameness prevention program requires an integrated strategy that addresses all the contributing factors:

- **Routine Hoof Care:** Implement a schedule for routine hoof trimming to maintain proper claw shape and balance. Trimming should be performed by a trained professional at least twice a year.
- **Optimal Housing and Environment:** Provide a clean, dry, and comfortable environment. Floors should be well-grooved to provide traction, and stalls should be designed to encourage sufficient lying time (12-14 hours per day) to reduce standing-related pressure on the hooves.
- **Footbathing:** Regular footbathing with disinfectants like copper sulphate or formalin helps to control infectious agents that cause conditions such as digital dermatitis.
- **Balanced Nutrition:** A diet that supports rumen health and avoids SARA is essential. Ensuring adequate intake of bypass fat and key trace minerals like zinc, copper, and biotin is also vital for maintaining a thick digital cushion and strong horn growth.
- **Monitoring and Early Detection:** Regular locomotion scoring and daily observation of the herd are crucial for identifying and treating lame cows at the earliest possible stage, which significantly improves recovery outcomes and prevents chronic issues.

Conclusion

Bovine lameness is a complex and costly condition, but with a deeper understanding of its aetiology—particularly the role of the digital cushion and the development of Claw Horn Disruption Lesions—we can develop more effective prevention and control strategies. By implementing an integrated management program that focuses on proactive hoof care, optimal housing and balanced nutrition, it is possible to significantly reduce the prevalence of lameness, thereby improving animal welfare and enhancing the long-term profitability and sustainability of a herd. The move from treating the symptom to addressing the root cause is the key to creating a healthier, more productive future for cattle.

Official Website
www.thescienceworld.net
thescienceworldmagazine@gmail.com
Published 30/03/2026

References

Manske, T., Hultgren, J., & Bergsten, C. (2002). The relationship between claw horn disruption lesions and locomotion score. *The Veterinary Journal*, *164*(2), 160–166.

Newsome, R., et al. (2017). The involvement of the digital cushion and the distal phalanx in the development and reoccurrence of claw horn disruption lesions in dairy cattle. University of Nottingham ePrints. Retrieved from https://eprints.nottingham.ac.uk/34436/