THE SCIENCE WORLD
A Monthly e Magazine
ISSN:2583-2212
Popular Article
Oct, 2026 Vol.6(10), 3388-3394

Dystocia In Ruminants And Companion Animals: When Birth Becomes An Emergency

M. Rajashri[1]* and K. Yashaswini[2]
[1]Assistant Professor, Department of Veterinary Gynaecology and Obstetrics, College of Veterinary Science, P.V. Narasimha Rao Telangana Veterinary University, Mamnoor, Warangal, Telangana, India. [2]Contract Teaching Faculty, Veterinary Clinical Complex, College of Veterinary Science, Mamnoor, Warangal, Telangana, India.
***e-mail:** rsri0835@gmail.com*
doi.org/10.5281/ScienceWorld.23959837

Introduction: The Critical Moments of Birth

Birth is one of the most remarkable events in an animal's life. After months of pregnancy, the arrival of a healthy newborn represents the successful completion of a complex biological process. Yet, sometimes, nature needs a helping hand. When an animal cannot deliver its offspring normally, the condition is known as **dystocia, or difficult birth**.

Dystocia is more than a problem during delivery. It is a veterinary emergency that may threaten the lives of both the mother and her unborn offspring. In livestock, difficult birth can lead to calf, lamb, or kid mortality, injuries to the mother, reduced milk production, delayed recovery, infertility, and significant economic losses. In companion animals such as dogs and cats, dystocia can result in the loss of valuable newborn puppies or kittens and, in severe cases, the death of the mother.

Although the underlying principles of birth are similar across animal species, the causes, warning signs, and methods of treatment differ considerably. Understanding these differences is essential for veterinarians, farmers, pet owners, and anyone involved in animal care.

Causes of Dystocia?

Broadly, the causes of dystocia are classified into two major categories:

Maternal causes: These originate from the mother and include weak or absent uterine contractions, incomplete cervical dilation, a narrow or injured pelvis, abnormalities of the reproductive tract, and uterine torsion, in which the uterus twists abnormally.

Fetal causes: These are associated with the unborn offspring and include excessive fetal size, abnormal posture, incorrect presentation, congenital abnormalities and, occasionally, the simultaneous entry of multiple fetuses into the birth canal. In many cases, both maternal and fetal factors contribute to the problem. Identifying the underlying cause is the first step towards choosing the safest method of delivery.

Dystocia in Ruminants: A Challenge for Livestock Production

Ruminants, including cattle, buffaloes, sheep and goats, are essential to agriculture and rural livelihoods. Their reproductive performance influences milk production, meat supply, farm profitability and household income. Consequently, difficult birth is not only a veterinary concern but also an important livestock-management issue.

1. Cattle and Buffaloes: When Calving Becomes Difficult

In cattle, one of the important causes of dystocia is **fetopelvic disproportion**, a mismatch between the size of the calf and the capacity of the mother's pelvic canal. This problem is particularly relevant in first-calving heifers, which may have a relatively small pelvic opening. Large calves, prolonged gestation, unsuitable sire selection, abnormal fetal posture, and failure of the cervix to dilate adequately can also contribute to difficult calving. Other important conditions include uterine torsion and uterine inertia, in which contractions are insufficient to expel the calf.

Buffaloes may experience similar reproductive difficulties. Because normal labour and the timing of intervention can vary among animals, careful observation and individual veterinary assessment are essential. The normal calf is usually delivered with its head resting between the two extended forelimbs. However, if the head is turned sideways, one or both forelimbs are flexed, or the hindquarters enter the birth canal in an abnormal posture, vaginal delivery may become difficult or impossible without assistance.

An important complication is uterine torsion. In this condition, the uterus rotates around its longitudinal axis, obstructing the birth canal and potentially compromising blood supply to reproductive tissues. The animal may show signs of discomfort, restlessness, reduced appetite, or failure to progress through labour. Prompt veterinary examination is essential because delayed treatment can endanger both the cow or buffalo and the fetus.

2. Sheep and Goats: Small Animals, Serious Emergencies

Dystocia in sheep and goats is often associated with abnormal fetal posture, multiple fetuses entering the birth canal together, incomplete cervical dilation, and inadequate uterine contractions. In goats, the presence of twins or triplets can occasionally complicate delivery when the fetuses are not correctly positioned. A head turned backwards, a flexed forelimb, or hindlimbs

Official Website
www.thescienceworld.net
thescienceworldmagazine@gmail.com
Published 30/10/2026

retained inside the birth canal may prevent normal delivery. In sheep, similar problems can occur, while fetal oversize and maternal pelvic limitations may also contribute.

One important condition in small ruminants is **ringwomb**, in which the cervix fails to dilate sufficiently for delivery. Unlike a simple malposition that may be corrected by an experienced Veterinarian, ringwomb can require more specialised intervention. Small ruminants are particularly vulnerable to injury from excessive traction. Their reproductive tissues can be damaged by forceful or repeated attempts to pull out a fetus. A seemingly minor delivery problem may therefore progress to vaginal injury, uterine rupture, infection or death.

Dystocia in Dogs and Cats: A Hidden Emergency at Home

Difficult birth is not limited to livestock. Dogs and cats may also experience serious complications during labour, sometimes when their owners are expecting a routine delivery at home. In companion animals, **uterine inertia**, or failure of the uterus to contract effectively, is a major cause of dystocia in dogs. It may occur from the beginning of labour or develop after one or more puppies have been delivered.

Other causes include oversized puppies or kittens, an abnormally narrow pelvic canal, fetal malposition, congenital abnormalities, and obstruction of the birth canal. Certain breeds are predisposed because of their body conformation. Brachycephalic dogs, including English Bulldogs and French Bulldogs, are particularly important examples because the relationship between fetal head size and the maternal pelvis can make natural delivery difficult. Cats can also experience dystocia, particularly when uterine contractions become inadequate, or a kitten is malpositioned. Although many queens deliver without assistance, prolonged labour should never be dismissed as normal simply because the animal appears calm between contractions.

Recognising the Warning Signs

Pet owners should seek urgent veterinary attention when:

- Strong, repeated abdominal contractions continue without producing a puppy or kitten.
- Labour appears to have stopped despite the presence of additional fetuses.
- A puppy or kitten is visibly lodged in the birth canal.
- The mother becomes extremely weak, collapses, develops severe pain or appears seriously unwell.
- There is profuse bleeding or abnormal, foul-smelling discharge.
- The expected delivery period has passed, and the animal shows signs of distress or labour fails to progress.

Green or greenish-black discharge before the first puppy or kitten is delivered can be a warning sign of placental separation, particularly in dogs, and warrants immediate veterinary advice. The

Official Website
www.thescienceworld.net
thescienceworldmagazine@gmail.com
Published 30/10/2026

timing of labour differs between species and individuals. Therefore, owners should not rely on a single time limit to decide whether a delivery is safe. The strength of contractions, progress of delivery, condition of the mother and evidence of fetal distress must all be considered.

Diagnosing a Dystocia?

The first task is to determine whether the problem arises from the mother, the fetus, or both. A veterinarian assesses the animal's general condition, pregnancy history, duration of labour, contraction pattern and progress of delivery. In ruminants, a carefully performed vaginal examination can help determine cervical dilation, fetal presentation, position and posture, and whether an obstruction is present. The examination must be conducted hygienically to reduce the risk of introducing infection or damaging reproductive tissues.

In dogs and cats, abdominal palpation, a vaginal examination when appropriate, radiography, and ultrasonography can help establish the number of fetuses, their position, approximate size, and viability. Ultrasonography is especially useful for assessing fetal heart activity and identifying signs of fetal compromise. The mother's condition is equally important. Dehydration, exhaustion, metabolic disturbances, shock, or reproductive-tract injury can influence the choice of treatment.

Treatment: Knowing When to Assist and When to Operate

There is no single treatment for every case of dystocia. The correct approach depends on the species, cause of the obstruction, duration of labour, fetal viability and the condition of the mother.

Assisted vaginal delivery

When the cervix is adequately dilated, the birth canal is unobstructed, and the fetus can be safely repositioned, an experienced veterinarian may correct abnormal presentation or posture and assist delivery using appropriate lubrication and controlled traction.

The objective is not simply to extract the fetus but to do so without causing injury to the mother or offspring. Excessive force, uncoordinated pulling and repeated unsuccessful attempts can transform a manageable case into a life-threatening emergency.

Medical management

Medicines that influence uterine contractions may be useful in selected cases of uterine inertia. However, they are not appropriate when the fetus is oversized, malpositioned or physically obstructed. Drugs such as oxytocin must be used only after veterinary assessment. When an obstruction is present, stimulating stronger uterine contractions can increase the risk of fetal distress, placental separation, uterine damage, or rupture.

Official Website
www.thescienceworld.net
thescienceworldmagazine@gmail.com
Published 30/10/2026

Calcium or other supportive treatments may be indicated in particular circumstances, depending on the species and the underlying clinical problem. These treatments require appropriate dosing and monitoring by a veterinarian.

Caesarean section

A caesarean section is a surgical procedure in which the fetus is delivered through an incision in the uterus. It may be necessary when vaginal delivery is impossible, the obstruction cannot be corrected safely, the mother is deteriorating, or the fetus is at risk because of prolonged labour. In cattle and buffaloes, surgery may be required in cases of severe fetopelvic disproportion, irreducible malpresentation or uterine torsion that cannot be corrected by other appropriate methods. In dogs and cats, a caesarean section may be lifesaving when medical management fails, or an obstruction makes vaginal delivery unsafe.

Although surgery involves cost, anaesthesia and postoperative care, delaying a necessary operation can increase the risk of infection, uterine injury, fetal death and maternal complications. The decision should be based on clinical findings rather than an attempt to avoid surgery at all costs.

The First Hours After Birth Matter Too

Successful delivery is only the beginning of neonatal care. Newborn animals must establish breathing, maintain body temperature, and receive appropriate nutrition.

In calves, lambs and kids, early intake of adequate colostrum is essential because it provides antibodies that help protect against infectious disease. The newborn should be checked for breathing difficulties, weakness, injuries and inability to suckle.

Puppies and kittens require warmth, adequate nursing and close observation. Newborns that are weak, cold, unable to suckle or struggling to breathe need prompt veterinary assessment. The mother also requires attention after dystocia. Examination for excessive bleeding, reproductive-tract injuries, retained fetal membranes where relevant, infection, and other postpartum complications helps protect her recovery and future fertility. Difficult delivery can be followed by reduced reproductive performance, so appropriate postpartum monitoring is important for both livestock and companion animals.

Prevention: The Best Treatment Begins Before Labour

Although not every case of dystocia can be prevented, many risk factors can be reduced through good reproductive management.

In livestock, prevention begins with appropriate breeding decisions. Selecting suitable sires, avoiding matings likely to produce excessively large calves, and ensuring that replacement heifers have achieved appropriate growth and pelvic development can reduce the risk of difficult calving.

Official Website
www.thescienceworld.net
thescienceworldmagazine@gmail.com
Published 30/10/2026

Balanced nutrition throughout pregnancy is equally important. Both undernutrition and excessive body condition can create problems. Mineral and nutritional management should be adapted to the animal's needs rather than relying on indiscriminate supplementation.

Pregnant animals should be observed more closely as delivery approaches. Farmers should maintain clean calving areas, keep essential veterinary contact details available and arrange early assistance when labour does not progress normally.

For dogs and cats, responsible breeding and pregnancy monitoring are key preventive measures. Breed-associated risks, previous dystocia, maternal age, general health and litter characteristics should be considered before breeding. Veterinary assessment during pregnancy can help establish expected delivery dates and identify animals that may require planned surgical delivery.

Owners should prepare a clean, quiet birthing area and know where emergency veterinary care is available. Importantly, they should never administer labour-inducing medicines or attempt forceful extraction of a newborn without veterinary guidance.

The Economic and Animal Welfare Dimensions

Dystocia has consequences that extend beyond the delivery room. In dairy and meat production systems, a difficult birth can increase veterinary expenses, prolong recovery, delay reproductive cycles, reduce production and result in the loss of valuable offspring.

The impact is particularly serious for small-scale farmers, for whom the loss of a single calf or buffalo may represent a substantial financial setback. The costs may also include extra labour, medicines, surgery and the treatment of postpartum complications.

In companion animals, dystocia can cause considerable emotional distress for owners and raise important questions about responsible breeding practices. Repeatedly breeding animals with severe conformational limitations without appropriate veterinary planning can place both mothers and offspring at avoidable risk.

From an animal-welfare perspective, timely intervention reduces prolonged pain, exhaustion, tissue injury and preventable mortality. It also reinforces the principle that breeding decisions must prioritise the health and well-being of the animals involved.

Conclusion: Timely Intervention Can Save Two Lives

Dystocia illustrates how a natural biological process can become a medical emergency when even one part of the birth process goes wrong. Whether it occurs in a cow in a rural dairy farm, a goat in a smallholder production system, or a family dog delivering puppies at home, the consequences can be severe if warning signs are ignored.

Official Website
www.thescienceworld.net
thescienceworldmagazine@gmail.com
Published 30/10/2026

The most important lessons are universal: recognise abnormal labour early, avoid forceful and untrained intervention, seek veterinary assistance promptly and ensure appropriate care for both the mother and her newborns.

With sound breeding practices, careful pregnancy monitoring, informed animal owners and timely veterinary treatment, many complications associated with difficult birth can be managed successfully. Protecting the mother and offspring is not merely a clinical responsibility; it is an essential part of sustainable livestock production, responsible pet ownership and compassionate animal care.

References

MSD Veterinary Manual. *Management of Calving in Cattle*. Merck & Co. Available at: https://www.msdvetmanual.com/management-and-nutrition/management-of-reproduction-cattle/management-of-calving-in-cattle

MSD Veterinary Manual. *Management of Dystocia in Cattle*. Merck & Co. Available at: https://www.merckvetmanual.com/management-and-nutrition/management-of-reproduction-cattle/management-of-dystocia-in-cattle

MSD Veterinary Manual. *Parturition in Goats*. Merck & Co. Available at: https://www.msdvetmanual.com/management-and-nutrition/management-of-reproduction-goats/parturition-in-goats

MSD Veterinary Manual. *Dystocia in Small Animals*. Merck & Co. Available at: https://www.merckvetmanual.com/reproductive-system/reproductive-diseases-of-the-female-small-animal/dystocia-in-small-animals

MSD Veterinary Manual. *Labor and Delivery in Dogs and Cats*. Merck & Co. Available at: https://www.msdvetmanual.com/management-and-nutrition/management-of-reproduction-dogs-and-cats/labor-and-delivery-in-dogs-and-cats